

On A Simple Construction Of Triangle Centers X(8), X(197), X(K) & X(594)

Dasari Naga Vijay Krishna
 Department of Mathematics, Narayana Educational Institutions
 Machilipatnam, Bangalore, INDIA
 E-mail: vijay9290009015@gmail.com

Abstract

In this article, we provide a new method for constructing the Nagel Point {X(8)}, Cevian quotient of Symmedian Point, Nagel Point {X(197)} and the isogonal conjugate of 1st Hatzipolakis-Yiu Point {X(594)}. In addition, we also establish some collinearity and concurrence.

Keywords: triangle centers, Nagel point, Carnot's theorem, collinearity and concurrence.

Introduction

In the literature of Encyclopedia of Triangle Centers [1], there is a list of over 2000 triangle centers. Among those X(8), X(197), X(K) and X(594) are four such triangle centers. In this note, we devote our study for the construction of these points and their related coincidence.

In article [2], Larry Hoehn gives another way to construct the Nagel point using only the incircle and not the excircles. In this article we deal with a simple and an elegant construction of these four points which reveals a new characterization of X(8), X(197), X(K), X(594) and their unexpected coincidence.

Terminology

Triangle center	Its name
X(8)	Nagel Point
X(197)	Cevian Quotient of Symmedian Point, Nagel Point
X(K)	Isogonal Conjugate of X(5019), Isotomic Conjugate of X(940), The Polar Conjugate of X(4185), The Trilinear Pole Of Line X(523)X(4391)
X(594)	Isogonal Conjugate of 1st Hatzipolakis-Yiu Point

Construction

Given a triangle ABC , let D, E, F be the midpoints of the sides BC, CA and AB , construct a circle O_A with A as center and AD as radius which intersects the line drawn through D and parallel to the internal angular bisector of angle A at L_A (it is clear that the point L_A is the reflection of D with respect to the external angular bisector of angle A). Similarly, define the points L_B and L_C . Consider the points N_A, N_B and N_C as

$$N_A = BL_C \cap CL_B, N_B = AL_C \cap CL_A \text{ and } N_C = BL_A \cap AL_B \text{ then}$$

(I) The line segments AN_A, BN_B and CN_C concur at X(594)

(II) The line segments $L_A N_A, L_B N_B$ and $L_C N_C$ concur at X(8)

Consider the points $N'_A, N'_B, N'_C, L'_A, L'_B, L'_C, M'_A, M'_B$ and M'_C as $N'_A = AN_A \cap BC$, $N'_B = BN_B \cap CA$, $N'_C = CN_C \cap AB$, $L'_A = AX(8) \cap BC$, $L'_B = BX(8) \cap CA$, $L'_C = CX(8) \cap AB$, $M'_A = L_A N_A \cap BC$, $M'_B = L_B N_B \cap CA$ and $M'_C = L_C N_C \cap AB$ then

(III) The line segments AM'_A, BM'_B and CM'_C are concur at $X(K)$

(IV) The lines $N_A L'_A, N_B L'_B$ and $N_C L'_C$ are concur at $X(197)$

(V) There exists a conic through any two triads out of three triads (D, E, F) ,

(N'_A, N'_B, N'_C) and (L'_A, L'_B, L'_C) .

In order to prove (1), (II), (III), (IV) and (V), we will make use of barycentric coordinates. If a triangle ABC has side lengths $BC = a$, $CA = b$, $AB = c$ then $A = (1 : 0 : 0)$, $B = (0 : 1 : 0)$, $C = (0 : 0 : 1)$, $D = (0 : 1 : 1)$, $E = (1 : 0 : 1)$ and $F = (1 : 1 : 0)$ in homogeneous barycentric coordinates with reference to ABC [4].

Preposition 1

The equation of line joining of two points with coordinates $(x_1 : y_1 : z_1)$ and

$$(x_2 : y_2 : z_2) \text{ is } \begin{vmatrix} x & y & z \\ x_1 & y_1 & z_1 \\ x_2 & y_2 & z_2 \end{vmatrix} = 0 \text{ or } x(y_1 z_2 - y_2 z_1) + y(z_1 x_2 - z_2 x_1) + z(x_1 y_2 - x_2 y_1) = 0$$

Preposition 2

The intersection of the two lines $p_1 x + q_1 y + r_1 z = 0$ and $p_2 x + q_2 y + r_2 z = 0$ is the point $(q_1 r_2 - q_2 r_1 : r_1 p_2 - r_2 p_1 : p_1 q_2 - p_2 q_1)$.

Preposition 3

Three lines $p_i x + q_i y + r_i z = 0$, $i = 1, 2, 3$, are concurrent if and only if

$$\begin{vmatrix} p_1 & q_1 & r_1 \\ p_2 & q_2 & r_2 \\ p_3 & q_3 & r_3 \end{vmatrix} = 0$$

Preposition 4

The barycentric coordinate of the points which are the reflection of an arbitrary point $P(u : v : w)$ with respect to the external angular bisector of angle A is

$$P_A = (c^2 v + b^2 w + b c (u + v + w) : -b^2 w : -c^2 v),$$

with respect to the external angular bisector of angle B is

$$P_B = (-a^2 w : c^2 u + a^2 w + a c (u + v + w) : -c^2 u),$$

and with respect to the external angular bisector of angle C is

$$P_C = (-a^2 v : -b^2 u : b^2 u + a^2 v + a b (u + v + w)) \text{ [4].}$$

Corollary

By replacing $u = 0$, $v = w = 1$ then $P_A = L_A = ((b+c)^2 : -b^2 : -c^2)$

By replacing $v = 0$, $u = w = 1$ then $P_B = L_B = (-a^2 : (a+c)^2 : -c^2)$

By replacing $w = 0$, $u = v = 1$ then $P_C = L_C = (-a^2 : -b^2 : (a+b)^2)$

Note

1. The triangles ABC and $P_AP_BP_C$ are perspective with perpector of the isogonal conjugate of P [4].
2. The excentral triangle and $P_AP_BP_C$ are perspective if and only if P lies on the Neuberg cubic of excentral triangle [4].

Using the prepositions listed above we list out the barycentric coordinates of the specified points and equation of the lines in barycentric system (we use standard notations such as $a = BC$, $b = CA$, $c = AB$ and $s =$ semi perimeter, $R =$ circum radius, $r =$ inradius, $\Delta =$ area).

Table 1. Point and its Barycentric Coordinates

Point	barycentric coordinates	Point	barycentric coordinates
L_A	$((b+c)^2 : -b^2 : -c^2)$	L'_A	$(0 : (s-b) : (s-c))$
L_B	$(-a^2 : (a+c)^2 : -c^2)$	L'_B	$((s-a) : 0 : (s-c))$
L_C	$(-a^2 : -b^2 : (a+b)^2)$	L'_C	$((s-a) : (s-b) : 0)$
N_A	$(-a^2 : (c+a)^2 : (a+b)^2)$	N'_A	$(0 : -(c+a)^2 : -(a+b)^2)$
N_B	$((b+c)^2 : -b^2 : (a+b)^2)$	N'_B	$(-(b+c)^2 : 0 : -(a+b)^2)$
N_C	$((b+c)^2 : (c+a)^2 : -c^2)$	N'_C	$(-(c+b)^2 : -(c+a)^2 : 0)$

Point	barycentric coordinates
M'_A	$(0 : sc^2+abc : sb^2+abc) = (0 : c^2+4Rr : b^2+4Rr)$
M'_B	$(c^2+4Rr : 0 : a^2+4Rr)$
M'_C	$(b^2+4Rr : a^2+4Rr : 0)$

Point	barycentric coordinates
$X(8)$	$((s-a) : (s-b) : (s-c)) = (\cot A/2 : \cot B/2 : \cot C/2)$
$X(197)$	$(a^2[s^2(b^2+c^2-a^2)-s(bc(b+c)+ca(c-a)+ab(b-a)+abc(b+c-a))] : b^2[s^2(c^2+a^2-b^2)-s(ca(c+a)+ab(a-b)+bc(c-b)+abc(c+a-b))] : c^2[s^2(a^2+b^2-c^2)-s(ab(a+b)+bc(b-c)+ca(a-c)+abc(a+b-c))])$ $= af(A,B,C) : bf(B,C,A) : cf(C,A,B)$ Where $f(A,B,C) = a[-a^2\tan A/2 + b^2\tan B/2 + c^2\tan C/2]$
$X(594)$	$((c+b)^2 : (c+a)^2 : (a+b)^2)$
$X(K)$	$((b^2+4Rr)(c^2+4Rr) : (c^2+4Rr)(a^2+4Rr) : (a^2+4Rr)(b^2+4Rr))$

Table 2. Line and its Barycentric Equation

Line	Equation	Line	Equation	Line	Equation
AL_B	$yc^2 + z(c+a)^2 = 0$	BL_A	$xc^2 + z(b+c)^2 = 0$	CL_B	$x(c+a)^2 + ya^2 = 0$
AL_C	$y(a+b)^2 + zb^2 = 0$	BL_C	$x(a+b)^2 + za^2 = 0$	CL_A	$xb^2 + y(b+c)^2 = 0$

Line	Equation
AN_A	$y(a+b)^2 - z(c+a)^2 = 0$
BN_B	$x(a+b)^2 - z(b+c)^2 = 0$
CN_C	$x(c+a)^2 - y(b+c)^2 = 0$

Line	Equation
$AX(8)$	$y(s-c) - z(s-b) = 0$
$BX(8)$	$x(s-c) - z(s-a) = 0$
$CX(8)$	$x(s-b) - y(s-a) = 0$

Line	Equation
$L_A N_A$	$xbc(b-c) - s[b^2(x+y) - c^2(x+z)] - abc(y-z) = 0$
$L_B N_B$	$yca(c-a) - s[c^2(y+z) - a^2(y+x)] - abc(z-x) = 0$
$L_C N_C$	$zab(a-b) - s[a^2(z+x) - b^2(z+y)] - abc(x-y) = 0$

Line	Equation
AM'_A	$y(b^2+4Rr) - z(c^2+4Rr) = 0$
BM'_B	$x(a^2+4Rr) - z(c^2+4Rr) = 0$
CM'_C	$x(a^2+4Rr) - y(b^2+4Rr) = 0$

Line	Equation
$N_A L'_A$	$a^2[y(s-c)-z(s-b)] + x(b-c)[s(b+c)-bc] = 0$
$N_B L'_B$	$b^2[z(s-a)-x(s-c)] + y(c-a)[s(c+a)-ca] = 0$
$N_C L'_C$	$c^2[x(s-b)-y(s-a)] + z(a-b)[s(a+b)-ab] = 0$

Theorem-1

The line segments AN_A , BN_B and CN_C are concurrent at $X(594)$.

Proof

Using table 2, it is easy to check that the point $X(594)\{((c+b)^2 : (c+a)^2 : (a+b)^2)\}$ satisfies the lines $AN_A \{y(a+b)^2 - z(c+a)^2 = 0\}$, $BN_B \{x(a+b)^2 - z(b+c)^2 = 0\}$ and $CN_C \{x(c+a)^2 - y(b+c)^2 = 0\}$.

Hence the lines AN_A , BN_B and CN_C are concurrent at $X(594)$. (see figure -1)




Figure 1.

Theorem-2

The line segments $L_A N_A$, $L_B N_B$ and $L_C N_C$ are concurrent at $X(8)$.

Proof

Using table 2, it is easy to check that the point $X(8)\{(s-a) : (s-b) : (s-c)\}$ satisfies the lines $L_A N_A$ $\{ xbc(b-c) - s[b^2(x+y) - c^2(x+z)] - abc(y-z) = 0 \}$, $L_B N_B$ $\{ yca(c-a) - s[c^2(y+z) - a^2(y+x)] - abc(z-x) = 0 \}$ and $L_C N_C$ $\{ zab(a-b) - s[a^2(z+x) - b^2(z+y)] - abc(x-y) = 0 \}$.

For example

Consider $xbc(b-c) - s[b^2(x+y) - c^2(x+z)] - abc(y-z) = 0 \dots\dots(1)$

Replace $x = (s-a)$, $y=(s-b)$, $z = (s-c)$, $x+y = c$, $x+z = b$ and $y-z = c-b$

$$\begin{aligned} xbc(b-c) - s[b^2(x+y) - c^2(x+z)] - abc(y-z) &= bc(s-a)(b-c) - s[b^2c - c^2b] - abc(c-b) \\ &= bc(s-a)(b-c) - bc(s-a)(b-c) = 0 \end{aligned}$$

Hence (1) is satisfied by the point $X(8)\{(s-a) : (s-b) : (s-c)\}$

That is the line $L_A N_A$ contains $X(8)$.

In the similar manner we can verify that $L_B N_B$ and $L_C N_C$ also contains $X(8)$.

So the lines $L_A N_A$, $L_B N_B$ and $L_C N_C$ are concurrent at **$X(8)$** . (see figure -2)

$$\begin{vmatrix} 0 & (b^2 + 4Rr) & -(c^2 + 4Rr) \\ (a^2 + 4Rr) & 0 & -(c^2 + 4Rr) \\ (a^2 + 4Rr) & -(b^2 + 4Rr) & 0 \end{vmatrix} = 0$$




Figure 3.

Note

The point X(K) is noted as the isogonal conjugate of X(5019) by *Francisco Javier* [5] and as the isotomic conjugate of X(940), the polar conjugate of X(4185) and the trilinear pole of line X(523), X(4391) by *Randy* [6].

Theorem 4

The lines $N_A L'_A$, $N_B L'_B$ and $N_C L'_C$ are concurrent at X(197)

Proof

Using table 2, it is easy to check that the point X(197) $\{af(A,B,C) : bf(B,C,A) : cf(C,A,B)\}$ satisfies the lines

$$N_A L'_A \{a^2[y(s-c)-z(s-b)] + x(b-c)[s(b+c)-bc] = 0\},$$

$$N_B L'_B \{b^2[z(s-a)-x(s-c)] + y(c-a)[s(c+a)-ca] = 0\}$$

$$\text{and } N_C L'_C \{c^2[x(s-b)-y(s-a)] + z(a-b)[s(a+b)-ab] = 0\}.$$

$$\text{Consider } a^2[y(s-c)-z(s-b)] + x(b-c)[s(b+c)-bc] = 0 \dots\dots(1)$$

$$\text{Replace } x = af(A, B, C), y = bf(B, C, A), z = cf(C, A, B)$$

$$\text{where } f(A, B, C) = a[-a^2 \tan A/2 + b^2 \tan B/2 + c^2 \tan C/2]$$

$$a^2[y(s-c)-z(s-b)] = a^2[b(s-c) f(B, C, A) - c(s-b) f(C, A, B)]$$

$$\begin{aligned}
 &= a^2 \{s[b f(B, C, A) - c f(C, A, B)] - bc [f(B,C,A) - f(C, A, B)] \} \\
 &= a^2 \{a^2 \tan A/2(b-c) (sb+sc-bc) - b^2 \tan B/2(b-c) (sb+sc-bc) - c^2 \tan C/2(b-c) (sb+sc-bc)\} \\
 &= -a^2(b-c) (sb+sc-bc) f(A, B, C) = -x(b-c)[s(b+c)-bc
 \end{aligned}$$

Hence (1) is satisfied by the point X(197) $\{af(A, B, C) : bf(B, C, A) : cf(C, A, B)\}$
That is the line $N_A L'_A$ contains X(197).

In a similar manner, we can verify that $N_B L'_B$ and $N_C L'_C$ also contains X(197).

So, the lines $N_A L'_A$, $N_B L'_B$ and $N_C L'_C$ are concur at X(197). (see figure-4)




Figure 4.

Preposition -5

Carnot's theorem

Suppose a conic L intersect in the side line BC at X, X' , CA at Y, Y' and AB at Z, Z' then

$$\left(\frac{BX}{XC}\right)\left(\frac{BX'}{X'C}\right)\left(\frac{CY}{YA}\right)\left(\frac{CY'}{Y'A}\right)\left(\frac{AZ}{ZB}\right)\left(\frac{AZ'}{Z'B}\right) = 1$$

Preposition -6

If X, Y, Z are the traces of a point P and X', Y' and Z' are the traces of another point Q on the sides of BC, CA , and AB of the triangle ABC such that $P = (u : v : w)$ and $Q = (u' : v' : w')$, then using Carnot's theorem there is a conic through six points. The equation of the conic is

$$\sum_{\text{cyclic}} \left[\frac{x^2}{uu'} - \left(\frac{1}{vw'} + \frac{1}{v'w} \right) yz \right] = 0, [4].$$

Theorem -5

a) There exists conic through (D, E, F) and (N'_A, N'_B, N'_C) whose barycentric equation is

$$\sum_{\text{cyclic}} \left[\frac{x^2}{(b+c)^2} - \left(\frac{1}{(a+b)^2} + \frac{1}{(c+a)^2} \right) yz \right] = 0$$

b) There exists conic through (N'_A, N'_B, N'_C) and (L'_A, L'_B, L'_C) whose barycentric equation is

$$\sum_{\text{cyclic}} \left[\frac{x^2}{(s-a)(b+c)^2} - \left(\frac{1}{(s-b)(a+b)^2} + \frac{1}{(s-c)(c+a)^2} \right) yz \right] = 0$$

c) There exists conic through (D, E, F) and (L'_A, L'_B, L'_C) whose barycentric equation is

$$\sum_{\text{cyclic}} \left[\frac{x^2}{(s-a)} - \left(\frac{1}{(s-c)} + \frac{1}{(s-b)} \right) yz \right] = 0$$

Proof:

Clearly the triads (D, E, F) , (N'_A, N'_B, N'_C) and (L'_A, L'_B, L'_C) are the traces of centroid $\{X(2)\}$ (1 : 1 : 1), the isogonal conjugate of first Hatzipolakis-Yiu Point $\{X(594)\}$ $((c+b)^2 : (c+a)^2 : (a+b)^2)$ and Nagel Point $\{X(8)\}$ $(s-a) : (s-b) : (s-c)$.

Hence, using Carnot's theorem there exists a conic through the traces of any two of the following three points $X(2)$, $X(594)$ and $X(8)$.

Their corresponding barycentric equations can be calculated using proposition 6.

Using tools similar to those used in this article, the following generalization can be demonstrated.

Generalization

In a given triangle ABC , $P_a(0 : v : w)$, $P_b(u : 0 : w)$, $P_c(u : v : 0)$ are the traces of an arbitrary point $P(u : v : w)$ on the sides BC, CA, AB respectively. P_A, P_B and P_C are the reflections of the points P_a, P_b, P_c with respect to the external angular bisectors of angles A, B and C . Let Q be the isogonal conjugate of P . Consider the points R_A, R_B and R_C as $R_A = BP_C \cap CP_B$, $R_B = AP_C \cap CP_A$ and $R_C = BP_A \cap AP_B$, then

(I) The line segments AR_A, BR_B and CR_C concur at R .

(II) The line segments P_AR_A, P_BR_B and P_CR_C concur at T .

Consider the points $R'_A, R'_B, R'_C, T'_A, T'_B, T'_C, M'_A, M'_B$ and M'_C as $R'_A = AR_A \cap BC$,

$$R'_B = BR_B \cap CA, R'_A = CR_A \cap AB, T'_A = AT \cap BC, T'_B = BT \cap CA, T'_C = CT \cap AB,$$

$$S'_A = P_AR_A \cap BC, S'_B = P_BR_B \cap CA \text{ and } S'_C = P_CR_C \cap AB \text{ then}$$

(III) The line segments AS'_A, BS'_B and CS'_C are concurrent at S .

(IV) The lines $R_A T'_A$, $R_B T'_B$ and $R_C T'_C$ are concur at Z .

(V) The points T , Q and R are collinear.

For further study about Nagel's point (X(8)) [2] is referred.

Acknowledgement

The author would like to thanks the anonymous referee for his/her kind comments and suggestions, which lead to a better presentation of this paper.

References

- [1] Kimberling C. *Encyclopedia of triangle centers*. Available from: <http://faculty.evansville.edu/ck6/encyclopedia/ETC.html>.
- [2] Krishna DNV. The fundamental property of Nagel point – A new proof. *J Sci Arts*. 2017; 1(38):31–36.
- [3] Hoehn L. A new characterization of the Nagel point. *Missouri J Math Sci*. 2007;19(1):45–48.
- [4] Yiu P. *Introduction to the Geometry of the Triangle* (Florida Atlantic University Lecture Notes: Summer 2011). Florida: Florida Atlantic University; 2013.
- [5] Francisco Javier. *Hyacinthos message 4293* (advanced plane geometry yahoo group), January, 2018.
- [6] Randy. *Hyacinthos message 4293* (advanced plane geometry yahoo group), January, 2018.